arXiv:0909.1128v2 [math.DG] 2 Aug 2010

APPLICATIONS OF A COMPLETENESS LEMMA IN MINIMAL SURFACE THEORY TO VARIOUS CLASSES OF SURFACES

M. UMEHARA AND K. YAMADA

Abstract. We give several applications of a lemma on completeness used by Osserman to show the meromorphicity of Weierstrass data for complete minimal surfaces with finite total curvature. Completeness and weak completeness are defined for several classes of surfaces which admit singular points. The completeness lemma is a useful machinery for the study of completeness in these classes of surfaces. In particular, we show that a constant mean curvature one (i.e. CMC-1) surface in de Sitter 3-space is complete if and only if it is weakly complete, the singular set is compact and all the ends are conformally equivalent to a punctured disk.

Introduction

Firstly, we recall the following lemma given in MacLane [10] and Osserman [17] to show properties of complete minimal surfaces. We set

$$\Delta := \{z \in \boldsymbol{C}\,;\,|z|<1\}, \qquad \text{and} \qquad \Delta^* := \Delta \setminus \{0\}.$$

Completeness Lemma ([10, 17], [18, Lemma 9.6])**.** *Let $\omega(z)$ be a holomorphic 1-form on Δ^*. Suppose that the integral $\int_\gamma |\omega|$ diverges to ∞ for all paths $\gamma : [0,1) \to \Delta^*$ accumulating at the origin $z=0$. Then $\omega(z)$ has at most a pole at the origin.*

This lemma plays a crucial role in showing the meromorphicity of Weierstrass data for complete minimal surfaces with finite total curvature. We show in the first section that it is also useful for showing several types of completeness of surfaces which may admit singularities.

In Section 2, we give a further application: Let $S^3_1(\subset \boldsymbol{R}^4_1)$ be the de Sitter 3-space with metric induced from the Lorentz-Minkowski 4-space $\boldsymbol{R}^4_1$, which is a simply-connected 3-dimensional complete Lorentzian manifold with constant sectional curvature 1. We consider the projection

$$p_L : \mathrm{SL}(2,\boldsymbol{C}) \longrightarrow S^3_1 = \mathrm{SL}(2,\boldsymbol{C})/\mathrm{SU}(1,1).$$

A holomorphic map $F\colon \Sigma^2 \to \mathrm{SL}(2,\boldsymbol{C})$ defined on a Riemann surface Σ^2 is called *null* if $\det(dF/dz)$ vanishes identically on Σ^2, where z is an arbitrary complex coordinate on Σ^2. For a null holomorphic immersion $F : \Sigma^2 \to \mathrm{SL}(2,\boldsymbol{C})$,

$$f := p_L \circ F : \Sigma^2 \longrightarrow S^3_1$$

gives a spacelike CMC-1 (i.e. constant mean curvature one) surface with singularities, called a *CMC-1 face*. Conversely, a CMC-1 face $f\colon \Sigma^2 \to \mathrm{SL}(2,\boldsymbol{C})$ is obtained as $f = p_L \circ F$, where $F\colon \widetilde{\Sigma}^2 \to \mathrm{SL}(2,\boldsymbol{C})$ is a null holomorphic immersion defined on the universal cover $\widetilde{\Sigma}^2$ of Σ^2 (see [1] and [2] for details). The pull-back metric $ds^2_\#$

Date: August 02, 2010.

2000 *Mathematics Subject Classification.* Primary 53C42; Secondary 53A35.

Key words and phrases. complete, CMC-1, mean curvature, Hopf differential, de Sitter space.

The authors are partially supported by Grant-in-Aid for Scientific Research (A) No. 19204005, and (B) No. 21340016, respectively, from the Japan Society for the Promotion of Science.

of the canonical Hermitian metric of $\mathrm{SL}(2,\boldsymbol{C})$ by the map F^{-1} of taking inverse matrices gives a single-valued positive definite metric on Σ^2 (see [2, Remark 1.11]). That is,

$$(1)\quad ds^2_{\#} := \operatorname{trace}(\alpha_{\#})^t\overline{(\alpha_{\#})} = \bigl(1+|G|^2\bigr)^2\left|\frac{Q}{dG}\right|^2 \qquad \Bigl(\alpha_{\#} := (F^{-1})^{-1}d(F^{-1})\Bigr),$$

where G and Q are the hyperbolic Gauss map and the Hopf differential, respectively (cf. [2, (1.17)]), and ${}^t(\ )$ denotes the transposition. A CMC-1 face $f:\Sigma^2 \longrightarrow S^3_1$ is called *weakly complete* if $ds^2_{\#}$ is complete.

On the other hand, we define another completeness as follows: Let (N^3,g) be a Riemannian or a Lorentzian manifold in general.

Definition 1. We say a C^∞-map $f\colon \Sigma^2 \to N^3$ is *complete* if there exists a symmetric covariant tensor T on Σ^2 with compact support such that ds^2+T gives a complete Riemannian metric on Σ^2, where ds^2 is the pull-back of the metric g on N^3, called the *first fundamental form.*

If f is complete and the singular set is non-empty, then the singular set must be compact, by definition. If f is a CMC-1 face, completeness implies weak completeness (cf. [2, Proposition 1.1]). This new definition of completeness is a generalization of the classical one, namely, if f has no singular points, then completeness coincides with the classical notion of completeness in Riemannian geometry. For complete CMC-1 faces, we have an analogue of the Osserman inequality for minimal surfaces in $\boldsymbol{R}^3$ (cf. [2, Theorem 0.2]). The goal of this paper is to prove the following assertion:

Theorem. *A CMC-1 face in S^3_1 is complete if and only if it is weakly complete, the singular set is compact and every end is conformally equivalent to a punctured disk.*

The ‘only if’ part has been proved in [2]. The proof of the ‘if’ part in this theorem is given in Section 2.

In this paper, we shall discuss the relationships between completeness and weak completeness on various classes of surfaces with singularities as applications of the completeness lemma. The above theorem is the deepest result amongst them.

1. Applications of the completeness lemma

In this section, we shall give several new applications of the completeness lemma, which show the importance of this kind of assertion (see Question in Remark 5).

Let (N^3,g) be a Riemannian 3-manifold. It is well-known that the unit tangent bundle T_1N^3 of N^3 has a canonical contact structure. A map $L:\Sigma^2\to T_1N^3$ defined on a 2-manifold Σ^2 is called a *Legendrian immersion* if and only if L is an isotropic immersion. Then a map $f:\Sigma^2\to N^3$ is called a *wave front* or a *front* if there exists a Legendrian immersion $L_f:\Sigma^2\to T_1N^3$ that is a lift of f. The pull-back metric

$$(2)\qquad d\tau^2 := L_f^*\tilde g$$

of the canonical metric $\tilde g$ of T_1N^3 coincides with the sum of the first fundamental form and the third fundamental form of f if (N^3,g) is a space form. Then f is called *weakly complete* if $L_f^*\tilde g$ is a complete Riemannian metric on Σ^2. It should be remarked that the completeness (in the sense of Melko and Sterling [14]) of surfaces of constant Gaussian curvature -1 in $\boldsymbol{R}^3$ coincides with our notion of weak completeness of wave fronts. The differential geometry of wave fronts is discussed in [19].

Remark 2. This weak completeness is different from that for CMC-1 faces as in the introduction. Moreover, weak completeness for improper affine spheres given in Remark 4 of this section is also somewhat different from that for fronts and CMC-1 faces. The authors do not yet know of a unified treatment of weak completeness. In fact, there might be several possibilities for completeness of a given class of surfaces with singularities.

Definition 1 defines completeness for wave fronts. By definition, completeness implies weak completeness. These two notions of completeness were defined for flat fronts (i.e. wave fronts whose first fundamental form has vanishing Gaussian curvature on their regular sets) in the hyperbolic 3-space H^3, the Euclidean 3-space $\boldsymbol{R}^3$, and the 3-sphere S^3, respectively (cf. [8], [15] and [6]). In particular, fundamental properties of flat surfaces in H^3 were given in Gálvez, Martínez and Milán [3]. Later, further properties for such surfaces as wave fronts were given in [7] and [8].

Let Σ^2 be a 2-manifold and $f:\Sigma^2\to H^3$ a flat front. In this case, there is another lifting of f (different from L_f) as follows: the map f induces a canonical complex structure on Σ^2, and there exists a holomorphic immersion $F:\widetilde{\Sigma}^2\to \mathrm{SL}(2,\boldsymbol{C})$ such that $f=\pi\circ F$ holds and $F^{-1}dF$ is off-diagonal, where

$$\pi:\mathrm{SL}(2,\boldsymbol{C})\longrightarrow H^3=\mathrm{SL}(2,\boldsymbol{C})/\mathrm{SU}(2)$$

is the canonical projection. Remarkably, the weak completeness of f is equivalent to the completeness of the pull-back metric of the canonical Hermitian metric of $\mathrm{SL}(2,\boldsymbol{C})$ by F (see [7]). For complete flat fronts in H^3, we have an analogue of the Osserman inequality for complete minimal surfaces (see [7] for details). The caustic (i.e. focal surface) of a complete flat front $f:\Sigma^2\to H^3$ is weakly complete (by taking a double cover of Σ^2 if necessary). The asymptotic behavior of weakly complete (but not complete) flat fronts in H^3 was analyzed in [9].

Proposition. *A flat front in H^3 is complete if and only if it is weakly complete, and the singular set is compact and each end of the front is conformally equivalent to a punctured disk.*

It should be remarked that another characterization of completeness of f using the total curvature of $d\tau^2$ is given in [8, Theorem 3.3].

Proof. The ‘only if’ part has been proved in [8, Theorem 3.3], since completeness at each end implies that the end is conformally equivalent to a punctured disk. So we shall prove the converse. We use the same notations as in [8]. Let $f:\Delta^*\to H^3$ be a flat immersion which is weakly complete at $z=0$. Then, the metric $d\tau^2$ given in (2) is written as $d\tau^2=|\omega|^2+|\theta|^2$, where ω and θ are the canonical forms as in [8, (2.4) and (2.7)]. Since f is weakly complete at $z=0$, the length $L(\gamma)$ with respect to the metric $d\tau^2=|\omega|^2+|\theta|^2$ diverges to ∞ for all paths $\gamma:[0,1)\to\Delta^*$ accumulating at the origin. Though ω is defined only on the universal cover $\widetilde{\Delta}^*$ of Δ^*, $|\omega|$ is well-defined on Δ^*. Hence there exists $\mu\in[0,1)$ such that

$$\omega(z)=z^{\mu}\hat{\omega}(z)\,dz, \tag{3}$$

where $\hat{\omega}(z)$ is a holomorphic function on Δ^*. (We have not yet excluded the possibility that $\hat{\omega}(z)$ has an essential singularity at $z=0$.) We shall now prove that $\hat{\omega}(z)$ has at most a pole there. For this purpose, we set $\rho:=\theta/\omega$. Then the singular set is characterized by $\{|\rho|=1\}$ (cf. [8, (2.6)]). We may assume $|\rho|<1$ on Δ^* without loss of generality by exchanging roles of ω and θ (see [8, page 270]), if necessary, because f is an immersion. Then we have that for any path γ accumulating at 0,

$$\infty=L(\gamma)=\int_\gamma\sqrt{|\omega|^2+|\theta|^2}\le\int_\gamma\sqrt{(1+|\rho|^2)|\omega|^2}\le\sqrt{2}\int_\gamma|\omega|\le\sqrt{2}\int_\gamma|\hat{\omega}(z)|\,|dz|.$$

Then by the completeness lemma, $\hat{\omega}(z)$ has at most a pole. On the other hand, $|\rho|$ is well-defined on Δ^* and $|\rho| < 1$ holds on Δ^*. Thus there exists a real number $\nu \in [0,1)$ such that $\rho(z) = z^{\nu}\hat{\rho}(z)$, where $\hat{\rho}(z)$ is a holomorphic function on Δ^* (that is, $\hat{\rho}(z)$ may have essential singularity at the origin). We have on Δ^* that

$$1 > |z^{1-\nu}| > |z^{1-\nu}\rho(z)| = |z\hat{\rho}(z)|.$$

Then by the great Picard theorem, $z\hat{\rho}(z)$ is meromorphic at the origin. Summing up, ω and $\theta = \rho\omega$ are meromorphic 1-forms on a neighborhood of the origin. Then [8, (3.2)] yields that $d\tau^2$ has finite total curvature, and completeness follows from [8, Theorem 3.3]. □

Remark 3. The assumption 'conformal equivalency of the end to the unit punctured disc' in Proposition cannot be dropped. However, in other situations, this condition might not be required. In fact, a flat front in $\boldsymbol{R}^3$ is complete if and only if it is weakly complete and the singular set is compact (see [15, Corollary 4.7]). On the other hand, a weakly complete immersion may not be complete in general. In fact, we set

$$ds^2 = du^2 + 2\cos\omega(u,v)\,du\,dv + dv^2$$

on $(\boldsymbol{R}^2; u, v)$, where $\omega : \boldsymbol{R}^2 \to (0,\pi)$ is a C^∞-function satisfying $\partial^2\omega(u,v)/\partial u\partial v = 0$. Then it is known that (cf [5] or [6]) there is a flat immersion $f : \boldsymbol{R}^2 \to S^3$ into the unit sphere S^3 such that ds^2 is the first fundamental form and $2\sin\omega(u,v)\,du\,dv$ is the second fundamental form. So if we set

$$\omega(u,v) = \arcsin\left(\frac{e^{-u^2}}{2}\right) + \arcsin\left(\frac{e^{-v^2}}{2}\right),$$

there is an associated flat immersion of $\boldsymbol{R}^2$ into S^3. In this case, the length of the curve $[0,\infty) \ni t \mapsto (t,-t) \in \boldsymbol{R}^2$ with respect to ds^2 is finite, that is

$$\int_0^\infty \sqrt{2\bigl(1-\cos\omega(t,-t)\bigr)} = \int_0^\infty 2\sin\frac{\omega(t,-t)}{2}dt = \int_0^\infty \exp\left(-t^2\right)dt < \infty.$$

Since the sum of the first and the third fundamental forms is given by $du^2 + dv^2$, the immersion f is weakly complete, but not complete.

Remark 4 (Improper affine spheres). Improper affine spheres in $\boldsymbol{R}^3$ are closely related to flat surfaces in H^3 (see [13] and also [4]). An improper affine sphere with admissible singular points in the affine space $\boldsymbol{R}^3$ is called an *improper affine map* (see [12]). Martínez [12] defined completeness for improper affine maps and proved an analogue of the Osserman inequality. Definition 1 is a generalization of this completeness and completeness for wave fronts. An improper affine map is called *weakly complete* if the metric $d\tau^2$ as in [12, (9)] is complete. Let Σ^2 be a Riemann surface and (F,G) a pair of holomorphic functions on Σ^2 such that

1. Re$(F\,dG)$ is exact,
2. $|dF|^2 + |dG|^2$ is positive definite.

Then the induced mapping $f : \Sigma^2 \to \boldsymbol{R}^3 = \boldsymbol{C} \times \boldsymbol{R}$ given by

$$f := \left(G + \overline{F}, \frac{|G|^2 - |F|^2}{2} + \operatorname{Re}\left(GF - 2\int F dG\right)\right)$$

is an improper affine map with $d\tau^2 = 2(|dF|^2 + |dG|^2)$. Conversely, any improper affine map is given in this way. The following assertion holds:

An improper affine map in $\boldsymbol{R}^3$ is a wave front. Moreover, it is complete if and only if it is weakly complete, the singular set is compact and all ends are conformally equivalent to a punctured disk.

In fact, $\nu := \bigl(\bar{F} - G, 1\bigr)$ gives a Euclidean normal vector field along f (cf. [12, (8)]), and one can easily verify that f is a wave front, that is, $(f, [\nu]) : \Sigma^2 \to \boldsymbol{R}^3 \times P^2(\boldsymbol{R})$ gives an immersion if and only if (ii) holds. (Nakajo [16, Proposition 4.3] gave an alternative proof of this fact.)

To prove the second assertion, we use the same notations as in [12]. Let $f : \Sigma^2 \to \boldsymbol{R}^3$ be a complete improper affine map. By [12, (9)], we have that

$$\begin{aligned} ds^2 &= |dF + dG|^2 \le (|dF| + |dG|)^2 \\ &= |dF|^2 + |dG|^2 + 2|dF||dG| \le 2(|dF|^2 + |dG|^2) = d\tau^2, \end{aligned}$$

which implies the completeness of $d\tau^2$, that is, f is weakly complete. On the other hand, by [12, Proposition 1], all ends of f are conformally equivalent to a punctured disk.

Next, we show the converse. Let $f : \Delta^* \to \boldsymbol{R}^3$ be an improper affine immersion which is weakly complete at $z = 0$, and for which the end $z = 0$ is of punctured type. Then the data (F, G) corresponding to f is a pair of holomorphic functions on Δ^*, and the length $L(\gamma)$ with respect to the metric $d\tau^2 = 2(|dF|^2 + |dG|^2)$ diverges to ∞ for all paths $\gamma : [0, 1) \to \Delta^*$ accumulating at the origin. Since f is an immersion, $|dF| \ne |dG|$ holds. So we may assume $|dF| < |dG|$ holds on Δ^* without loss of generality. Then we have that

$$\infty = L(\gamma) = \sqrt{2} \int_\gamma \sqrt{|dF|^2 + |dG|^2} \le 2 \int_\gamma |dG|.$$

Thus, by the completeness lemma, $dG(z)$ has at most a pole at the origin. Since $|dF/dG| < 1$, the great Picard theorem yields that $dF(z)/dG(z)$ has at most a pole at the origin. In particular, both dF and dG have at most a pole at the origin. Thus $dF(0)/dG(0)$ is well-defined, and satisfies $|dF(0)/G(0)| \le 1$. Since the singular set $\{z \in \Delta^* \,;\, |dF(z)/dG(z)| = 1\}$ is empty, we have that $|dF(0)/dG(0)| < 1$. Then there exists $\delta \in (0, 1)$ such that $|dF(z)/dG(z)| < \delta$ holds near $z = 0$, and

$$\begin{aligned} ds^2 = |dG + dF|^2 = |dG|^2 \left| 1 + \frac{dF}{dG} \right|^2 &\ge |dG|^2 \left(1 - \left| \frac{dF}{dG} \right| \right)^2 \\ \ge (1 - \delta)^2 |dG|^2 &\ge \frac{(1 - \delta)^2}{2} \left(|dF|^2 + |dG|^2 \right), \end{aligned}$$

which proves the completeness of f.

2. Proof of Theorem

Now we give a proof of the theorem in the introduction. It is sufficient to show the converse of [2, Proposition 1.1]. We use the same notations as in [2].

Let $f \colon \Delta^* \to S^3_1$ be a CMC-1 immersion which is weakly complete at the origin. Since f is an immersion, the metric $d\sigma^2$ given in [2, (1.15)] is a metric of constant curvature -1 on Δ^*. By [2, Theorem 2.1 and Definition 3.2], f is g-regular. Also, by [2, Corollary 3.1], f must be elliptic or parabolic.

Elliptic case. In this case, f is a g-regular elliptic end. Since the Schwarzian derivative $S(g)$ of the secondary Gauss map g is a projective connection of elliptic type (see [2, Section 2]), g is written in the form (see [2, Proposition 2.2])

$$g = z^\mu h(z),$$

where μ is a real number and h is a holomorphic function with $h(0) \ne 0$. If $\mu \ne 0$, then we can replace g by $1/g$ (see [2, Remark 1.9]), and so we may assume that $\mu > 0$. In particular, $g(0) = 0$ holds. On the other hand, if $\mu = 0$, then $|g(0)| \ne 1$, since the singular set $\{|g| = 1\}$ does not accumulate at the origin. Then, replacing g by $1/g$ if necessary (cf. [2, Remark 1.9]), we may assume $|g|^2 < 1 - \varepsilon$ on Δ^* for sufficiently small $\varepsilon > 0$. When $\mu > 0$, the inequality $|g|^2 < 1-\varepsilon$ holds trivially. Thus we have that

$$d\hat{s}^2 := (1+|g|^2)^2|\omega|^2 \le 4|\omega|^2 \le \frac{4}{\varepsilon^2}(1-|g|^2)^2|\omega|^2 = \frac{4}{\varepsilon^2}ds^2, \tag{4}$$

where (g,ω) is the Weierstrass data as in [2, (1.6)].

The weak completeness means the completeness of the metric $ds^2_{\#}$ as in (1). (The metric $ds^2_{\#}$ coincides with the metric given in [2, (1.17)].) On the other hand, the completeness of $ds^2_{\#}$ is equivalent to that of $d\hat{s}^2$ (see the last part of the proof of [2, Proposition 1.1]). By (4), ds^2 is complete at the origin.

Parabolic case. In this case, by [2, Lemma P], f is a g-regular parabolic end of the first kind. In the proof of Theorem 3.2 in [2], completeness is not used, but what is applied is the fact that the ends are immersed, g-regular and of punctured type. So we can refer to all of equations in that proof. One can choose the secondary Gauss map g as (see [2, (3.2)])

$$\frac{1}{i}\frac{g(z)+1}{g(z)-1} = \hat{g}(z) = i(h(z)+\varepsilon\log z),$$

where $\varepsilon = \pm 1$ and $h(z)$ is a holomorphic function on a sufficiently small neighborhood of the origin. Then we have that

$$g(z) := \frac{\hat{g}(z)-i}{\hat{g}(z)+i}, \qquad g'(z) = \frac{2(zh'(z)+\varepsilon)}{z(h(z)+\varepsilon\log z+1)^2}, \qquad \left({}' = \frac{d}{dz}\right).$$

Since $zh'(z)+\varepsilon$ is bounded on a neighborhood of the origin and

$$\frac{|(h(z)+\varepsilon\log z+1)|}{|\log z|} \to 1 \tag{5}$$

as $z \to 0$, there exists a positive constant c_1 such that

$$|g'| \le \frac{c_1}{|z|\,|\log z|^2}. \tag{6}$$

On the other hand, by (5) we have that

$$|g(z)| = \left|\frac{h(z)+\varepsilon\log z-1}{h(z)+\varepsilon\log z+1}\right| \to 1 \qquad (z\to 0). \tag{7}$$

Moreover, since

$$1-|g|^2 = \frac{4(\mathrm{Re}h+\varepsilon\log|z|)}{|\log z+\varepsilon(h+1)|^2},$$

it can be easily checked that there is a constant c_2 such that

$$\frac{c_2}{|\log z|} \le |1-|g|^2|. \tag{8}$$

Since $ds^2 = (1-|g|^2)^2|Q/dg|^2$, (6) yields that there is a constant $c_2'(>0)$ such that

$$|z|^2|\log z|^2\left|\frac{Q}{dz}\right|^2 \le c_2' ds^2, \tag{9}$$

where $Q = \omega\, dg$ is the Hopf differential (cf. [2, (1.8)]). Then, there exist positive constants c_3 and c_3' such that

$$d\hat{s}^2 := (1+|g|^2)^2|\omega|^2 \le c_3\left|\frac{Q}{dg}\right|^2 \le c_3'\,|\log z|^4\left|\frac{Q}{dz}\right|^2. \tag{10}$$

We recall the following inequality shown on [2, Appendix A]

$$ds^2 \le c_0|Q/dz|^2, \tag{11}$$

which holds for g-regular parabolic ends of the first kind, where c_0 is a positive constant. By (8), (9) (10) and (11), we have that

$$d\hat{s}^2 \le c_3' \left|\log z\right|^4 \left|\frac{Q}{dz}\right|^2 \le c_3' c_2' \frac{|\log z|^2}{|z|^2} ds^2 \le c_3' c_2' c_0 \frac{|\log z|^2}{|z|^2} \left|\frac{Q}{dz}\right|^2$$
$$\le \frac{c_3'(c_2')^2 c_0}{|z|^4} ds^2 \le c_3'(c_2' c_0)^2 \left|\frac{Q}{z^2 dz}\right|^2 .$$

Again, by the equivalency of completeness of the two metrics $ds^2_{\#}$ and $d\hat{s}^2$, the weak completeness implies the completeness of $d\hat{s}^2$. In particular, $|Q/(z^2dz)|^2$ is a complete metric at the origin. Then by the completeness lemma, $Q/(z^2dz)$ (and so Q) has at most a pole at $z = 0$. We denote by $\mathrm{ord}_0 Q$ the *order* of Q at the origin. For example, $\mathrm{ord}_0 Q = m$ if $Q = z^m\, dz^2$.

Firstly, we consider the case that $z = 0$ is not a regular end. Then the hyperbolic Gauss map $G(z)$ of f has an essential singular point at $z = 0$ and the Schwarzian derivative $S(G)$ has pole of order < -2. Since f is g-regular, $S(g)$ is of order -2 (see [2, Definition 3.2]), so the identity $2Q = S(g) - S(G)$ (cf. [2, (1.20)]) implies $\mathrm{ord}_0 Q \le -3$. On the other hand, if $z = 0$ is a regular end, then $\mathrm{ord}_0 Q = -2$ by [2, Lemma 5.1]. Thus, the Hopf differential of f satisfies $\mathrm{ord}_0 Q \le -2$. In particular, there exists a constant $c_4 > 0$ such that

$$|Q| \ge \frac{c_4 |dz|^2}{r^2} \qquad (r := |z|)$$

holds on Δ^*. By [2, (3.4)], we have that

$$d\sigma^2 := \frac{4\,|dg|^2}{(1-|g|^2)^2} \le \frac{C^2|dz|^2}{r^2(c+\log r)^2},$$

where C and c are positive constants. Thus there exists a constant $c_5 > 0$ such that (cf. [2, (1.15) and (1.16)])

$$ds^2 = 4\frac{|Q|^2}{d\sigma^2} \ge 4(c_5)^2 r^2 |Q|^2 (c+\log r)^2 \ge 4(c_4c_5)^2 \frac{(c+\log r)^2}{r^2} |dz|^2.$$

Since $|dz| \ge dr$ and

$$\int_1^t \frac{(c+\log r)}{r} dr = \frac{(\log t)^2}{2} + c\log t$$

diverges to ∞ as $t \to 0$, the metric ds^2 is complete at $z = 0$, which proves the assertion.

Remark 5. Related to the above proof of our main theorem, we leave here the following:

Question. Let $\omega(z)$ be a holomorphic 1-form on Δ^* and n a non-negative integer. Suppose that the integral

$$\int_\gamma |\omega(z)(\log z)^n|$$

diverges to ∞ for all C^∞-paths $\gamma : [0,1) \to \Delta^*$ accumulating at the origin $z = 0$. Then does $\omega(z)$ have at most a pole at the origin?

When $n = 0$, this question reduces to the original lemma. If the answer is affirmative, one can obtain the meromorphicity of the Hopf differential Q directly by applying it to equation (10), since the weak completeness implies the completeness of the metric $d\hat{s}^2$. The proof of the completeness lemma given in [18] cannot be modified directly, since the estimate of $\mathrm{Im}(\log z)$ along a path γ seems difficult. Fortunately, in our situation, we have succeeded to prove our main result without applying the statement, because of (11). It should be remarked that this key inequality (11) itself comes from the Gauss equation of the surface.

Remark 6 (Spacelike maximal surfaces in $\boldsymbol{R}^3_1$). It is known that CMC-1 surfaces in S^3_1 have a quite similar properties to spacelike maximal surfaces in the Lorentz-Minkowski space $\boldsymbol{R}^3_1$ of dimension 3 with the metric of signature $(-,+,+)$. We consider a fibration

$$p_L : \boldsymbol{C}^3 \ni (\zeta^1, \zeta^2, \zeta^3) \longmapsto \operatorname{Re}\left(-\sqrt{-1}\zeta^3, \zeta^1, \zeta^2\right) \in \boldsymbol{R}^3_1.$$

The projection of null holomorphic immersions into $\boldsymbol{R}^3_1$ by p_L gives spacelike maximal surfaces with singularities, called *maxfaces* (see [20] for details). Here a holomorphic map $F = (F_1, F_2, F_3) \colon \Sigma^2 \to \boldsymbol{C}^3$ defined on a Riemann surface Σ^2 is called *null* if $(F_1')^2 + (F_2')^2 + (F_3')^2$ vanishes identically, where $' = d/dz$ denotes the derivative with respect to a local complex coordinate z of Σ^2. The completeness and the weak completeness for maxfaces are defined in [20]. As in the case of CMC-1 faces in S^3_1, completeness implies weak completeness, however, the converse is not true. For complete maxfaces, we have an analogue of the Osserman inequality (cf. [20]). On the other hand, recently, the existence of many weakly complete bounded maxfaces in $\boldsymbol{R}^3_1$ has been shown, and such surfaces cannot be complete (see [11]). We can prove the following assertion by applying the completeness lemma:

A maxface in $\boldsymbol{R}^3_1$ is complete if and only if it is weakly complete, the singular set is compact and all ends are conformally equivalent to a punctured disk.

The proof of this assertion is easier than the case of CMC-1 faces in S^3_1. The 'only if' part has been proved in [20, Corollary 4.8], since finiteness of total curvature implies that all ends are conformally equivalent to a punctured disk. So it is sufficient to show the converse. The notations are the same as in [20]. Let $f : \Delta^* \to \boldsymbol{R}^3_1$ be a spacelike maximal immersion which is weakly complete at $z = 0$, namely, the length $L(\gamma)$ with respect to the metric $d\sigma^2 = (1 + |g|^2)^2|\omega|^2$ given in [20, Definition 2.7] diverges to ∞ for all paths $\gamma : [0, 1) \to \Delta^*$ accumulating at the origin. Since f is an immersion, we may assume that $|g(z)| \neq 1$ for all $z \in \Delta^*$. In particular, the image $g(\Delta^*)$ has infinitely many exceptional values. Then by the great Picard theorem, g has at most a pole at the origin. Without loss of generality, we may assume that $g(0) \in \boldsymbol{C}$. Then there exists $M > 0$ such that $|g(z)| < M$ for $z \in \Delta^*$. Thus we have that

$$\infty = L(\gamma) = \int_\gamma d\sigma \leq (1 + M^2) \int_\gamma |\omega|.$$

By the completeness lemma, ω has at most a pole at the origin. Since the Weierstrass data (g, ω) has at most a pole at the origin, $d\sigma^2$ has finite total curvature. Then completeness follows from [20, Corollary 4.8].

References

[1] S. Fujimori , 'Spacelike CMC 1 surfaces with elliptic ends in de Sitter 3-space' *Hokkaido Math. J.* 35 (2006) 289–320.

[2] S. Fujimori, W. Rossman, M. Umehara, K. Yamada and S.-D. Yang , 'Spacelike mean curvature one surfaces in de Sitter 3-space', *Comm. Anal. Geom.* 17 (2009) 383–427.

[3] J. A. Gálvez, A. Martínez and F. Milán , 'Flat surfaces in hyperbolic 3-space', *Math. Ann.* 316 (2000) 419–435.

[4] G. Ishikawa and Y. Machida , 'Singularities of improper affine spheres and surfaces of constant Gaussian curvature', *Int. J. Math.* 17 (2006) 269–293.

[5] Y. Kitagawa , 'Periodicity of the asymptotic curves on flat tori in S^3', *J. Math. Soc. Japan* 40 (1988) 457–476.

[6] Y. Kitagawa and M. Umehara , 'Extrinsic diameter of immersed flat tori in S^3', preprint.

[7] M. Kokubu, M. Umehara and K. Yamada , 'Flat fronts in hyperbolic 3-space', *Pacific J. Math.* 216 (2004) 149–175.

[8] M. Kokubu, W. Rossman, M. Umehara and K. Yamada , 'Flat fronts in hyperbolic 3-space and their caustics', *J. Math. Soc. Japan* 59 (2007) 265–299,

[9] M. Kokubu, W. Rossman, M. Umehara and K. Yamada , 'Asymptotic behavior of flat surfaces in hyperbolic 3-space', *J. Math. Soc. Japan* 61 (2009) 799–852.

[10] G. R. MacLane , 'On asymptotic values', abstract 603-166, *Notice of the Amer. Math. Soc* 10 (1963) 482–483.

[11] F. Martín, M. Umehara and K. Yamada , 'Complete bounded holomorphic curves immersed in $\boldsymbol{C}^2$ with arbitrary genus', *Proc. Amer. Math. Soc.* 137 (2009), 3437–3450.

[12] A. Martínez , 'Improper Affine maps', *Math. Z.* 249 (2005) 755–766,

[13] A. Martínez , 'Relatives of Flat Surfaces in H^3', Proceedings of INTERNATIONAL WORKSHOP ON INTEGRABLE SYSTEMS, GEOMETRY AND VISUALIZATION (November 19–24 at Kyushu University, Fukuoka, Japan), pp. 115–132, 2005.

[14] M. Melko and I. Sterling , 'Application of soliton theory to the construction of pseudospherical surfaces in $\boldsymbol{R}^3$', *Annals of Global Analysis and Geometry* 11 (1993) 65–107.

[15] S. Murata and M. Umehara , 'Flat surfaces with singularities in Euclidean 3-space', *J. of Differential Geometry* 82 (2009) 279–316.

[16] D. Nakajo, 'A representation formula for indefinite improper affine spheres', *Result. Math.* 55 (2009), 139–159.

[17] R. Osserman , 'Global Properties of Minimal Surfaces in E^3 and E^n', *Ann. of Math.* 80 (1964), 340–364.

[18] R. Osserman , *A survey of minimal surfaces* (Dover Publications Inc, 1986).

[19] K. Saji, M. Umehara and K. Yamada , 'The geometry of fronts', *Ann. of Math.* 169 (2009) 491–529.

[20] M. Umehara and K. Yamada , 'Maximal surfaces with singularities in Minkowski space', *Hokkaido Math. J.* 35 (2006) 13–40.

(Masaaki Umehara) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE,, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN
E-mail address: `umehara@math.sci.osaka-u.ac.jp`

(Kotaro Yamada) DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, O-OKAYAMA, MEGURO, TOKYO 152-8551, JAPAN
E-mail address: `kotaro@math.titech.ac.jp`